# Florida Florida FLUREVIEW Summer 2012



# Week 21: May 20-26, 2012



## Summary

The Florida Department of Health (FDOH) monitors multiple surveillance systems such as the Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE), the Florida Pneumonia and Influenza Mortality Surveillance System (FPIMSS), notifiable disease reports (Merlin), EpiCom, and Florida ILINet in order to track influenza activity in the state.

State:

- Influenza and ILI activity in Florida is low in most FDOH surveillance systems. ESSENCE Emergency
  Department data show elevated percent ILI in some counties, compared to previous years at this
  time. Most counties report low to no activity. Three counties report moderate activity.
- No ILI or influenza outbreaks were reported to EpiCom in week 21.
- In week 21, 8 specimens tested PCR-positive for influenza at the state lab. Seven specimens tested
  positive for influenza B, one specimen tested positive for influenza A unspecified. In recent weeks,
  influenza B has been the most common strain identified by the state lab. Other viruses known to be
  currently circulating, potentially causing ILI, include adenovirus, rhinovirus, parainfluenza, and
  respiratory syncytial virus (RSV).

TABLE 1: Summary of Florida Influenza-Like Illness (ILI) Activity for Week 21

Measure	Difference from previous week	Current week 21	Previous week 20	Page of Report
Percent of visits to ILINet providers for ILI	▼ 1.6	3.0%	1.4%	2
Percent of emergency department visits (from ESSENCE) due to ILI	▲ 0.1	2.6%	2.5%	3
Percent of laboratory specimens that were positive for influenza	▼ 9.5	22.8%	32.3%	5
Number of counties reporting moderate influenza activity	<b>▲</b> 1	3	2	6
Number of counties reporting widespread influenza activity	No Change	0	0	6
Number of counties reporting increasing influenza activity	<b>A</b> 4	7	3	7
Number of counties reporting decreasing influenza activity	▼ 12	4	16	7
Number of ILI outbreaks reported in EpiCom	No Change	0	0	10

### May 30, 2012

Posted on the Bureau of Epidemiology website: <a href="http://www.doh.state.fl.us/floridaflu/">http://www.doh.state.fl.us/floridaflu/</a>

Produced by: Bureau of Epidemiology, Florida Department of Health (FDOH)

Contributors: Heather Rubino, MS; Colin Malone, MPH; Aaron Kite-Powell, MS; Leah Eisenstein, MPH; Lillian Stark, PhD, MPH, MS; Valerie Mock; Janet Hamilton, MPH; Richard Hopkins, MD, MSPH

### In this Issue: Summary 1 **NREVSS Respiratory Virus** 1 Surveillance Outpatient Influenza-like Illness Surveillance Network (ILINET)-Statewide Influenza and ILI outbreaks reported 2 in EpiCom **ESSENCE Syndromic Surveillance** Summary-Statewide **ESSENCE Syndromic Surveillance** Summary-Regional Florida Bureau of Laboratories Viral Surveillance County Influenza Activity Map 6

**County Influenza Activity Trend** 

7


# NREVSS Respiratory Virus Surveillance

The National Respiratory and Enteric Virus Surveillance System (NREVSS) collects data from laboratory facilities around the country on a weekly basis. NREVSS monitors temporal and geographic patterns of RSV, human parainfluenza viruses, human metapneumo virus (HMPV), respiratory and enteric adenoviruses, and rotavirus. Florida has over 30 participating laboratory facilities.

FIGURE 1 shows the percentage of positive tests for multiple respiratory viruses reported by NREVSS-participating laboratories in Florida

The six respiratory viruses summarized in Figure 1 are:

- RSV
- Parainfluenza 1-3
- Adenovirus
- HMPV
- Rhinovirus
- Influenza



ILINet is a nationwide surveillance system composed of sentinel providers. Florida has 110 providers enrolled in ILINet who submit weekly ILI and total visit counts, as well as submitting ILI specimens to the BOL for confirmatory testing.

**FIGURE 2** shows the percentage of visits for ILI\* reported by ILINet Sentinel Providers statewide.


ILI percent positive remains low and similar to other non-pandemic seasons at this time. ILINet Provider reporting declines in the summer months. Figures will be updated as new data are received.

49 of 110 ILINet Sentinels have reported visit counts as of 11:00 a.m., May 30, 2012.


9 of 16 ILINet Super-Sentinels have reported visit counts as of 11:00 a.m., May 30, 2012.

**FIGURE 3** shows ILI visit counts reported by ILINet sentinel providers statewide by age group.

ILI visit counts are decreasing in the 0-55+ age group in week 21.



\*ILI = Influenza-like illness, fever >100°F AND sor e throat and/or cough *in the absence* of another known cause. ^There is no week 53 during the 2009-2010, 2010-2011, and 2011-12 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1.



†Data presented here are counts, not proportions as included in Figure 2. This is because age group denominator data is not available through ILINet.

# **Influenza and ILI Outbreaks**


In week 21 there were no influenza or ILI outbreaks reported into EpiCom.

Eleven influenza or ILI outbreaks were reported into EpiCom in the 2011-12 influenza season.

Florida uses ESSENCE for syndromic surveillance, which currently collects data daily from 174 hospital emergency departments (ED). These data are processed into 11 different syndrome categories based on the patient's chief complaint. One of the categories is ILI, which is composed of chief complaints that include the words "influenza" or "flu," or complaints that contain fever plus cough and/or sore throat.

**FIGURE 4** shows ESSENCE data on ILI visits to EDs as a percentage of all ED Visits.


Overall activity for influenza-like illness reported in ESSENCE is elevated compared to levels seen in previous non-pandemic seasons at this time.



\*There is no week 53 for the 2007-2008, 2009-2010, and 2010-2011 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1.

**FIGURE 5** shows percentage of ILI among all ED visits by age group.


Age-specific trends show that percent ILI in the 0-55+ age group is increasing in week 21.



One hundred five ESSENCE participating facilities are able to provide discharge disposition data for their ED visits going back to week 40, 2010. Using this information, the percent of ED visits for ILI that result in hospitalization can be calculated. The highest percentage of admissions is in the 55+ years old age group. The low number of visits in the 55+ age group causes variability in the ILI admission percentage from week to week.

**FIGURE 6** shows the percentage of ED visits for ILI that resulted in hospitalization, by age group.

Figure 6: Percentage of ED Visits Resulting for ILI Resulting in Hospital Admission, Hospitals Reporting Discharge Disposition (N=105), Week 40, 2010 to Week 21, 2012



### FIGURE 7 - FIGURE 13 describe ED chief complaint data from ESSENCE by Regional Domestic Security Task Force (RDSTF).

ILI activity in ESSENCE is elevated over previous years at this time in Regions 3, 5, 6 and 7.

Map 1: Hospitals Reporting Emergency Department (ED) Data to Florida ESSENCE, May 30, 2012 (N=174)







Figure 8: Percentage of Influenza-Like Illness from Emergency Department (ED) Chief  $Complaints, RDSTF\ Region\ 2\ ESSENCE\ Participating\ Hospitals\ (N=2),\ Week\ 1,\ 2009$ through May 30, 2012










Figure 9: Percentage of Influenza-Like Illness from Emergency Department (ED) Chief Complaints, RDSTF Region 3 ESSENCE Participating Hospitals (N=14), Week 40, 2008 through May 30, 2012




Figure 11: Percentage of Influenza-Like Illness from Emergency Department (ED) Chief Complaints, RDSTF Region 5 ESSENCE Participating Hospitals (N=49), Week 40, 2008 through May 30, 2012





Figure 13: Percentage of Influenza-Like Illness from Emergency Department (ED) Chief Complaints, RDSTF Region 7 ESSENCE Participating Hospitals (N=48), Week 40, 2008



<sup>\*</sup>There is no week 53 for the 2009-2010, 2010-2011 and 2011-2012 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1.

<sup>\*\*</sup>Historical data for region 2 is only available beginning week 1, 2009

**Table 2** shows the number of specimens tested by the Bureau of Laboratories (BOL), how many are influenza positive, and how many are H1N1 or other influenza subtypes.

**FIGURE 14 - FIGURE 15** use BOL viral surveillance data to track the progress of influenza infection over time. They include weekly information on how many specimens are tested by the BOL, what proportion of those test positive for influenza, and what subtypes are found for the positive influenza specimens.


Small numbers of influenza specimens submitted to BOL tested positive for influenza A H3, 2009 H1N1, and influenza B.


Influenza B has been the most common strain detected by BOL in recent weeks.

Table 2: Bureau of Laboratories Viral Surveillance for Week 21 by Lab Event Date\* as reported by 1:30 p.m. May 30, 2012

	Current Week 21	Previous Week 20
Total Specimens Tested	35	37
Influenza Positive Specimens (% of total)	8 (22.8%)	12 (32.3%)
H1N1 Positive Specimens (% of influenza positives)	-	1 (8.3%)
H3 Influenza A	-	3 (25 %)
Influenza A Unspecified	1 (12.5%)	-
Influenza B Unspecified	7 (77.1 %)	8 (66.7 %)

Figure 14: Number of Influenza-Positive Specimens Tested by the Florida Bureau of Laboratories (BOL) by Subtype by Lab Event Date\* Week 1, 2011 to Week 21, 2012 as Reported in Merlin by 11:30 a.m. May 30, 2012





For county-specific laboratory data, please refer to the Flu Lab Report in Merlin.

For instructions on how to use the Flu Lab Report, please see the Guide to Flu Lab Report on the Bureau of Epidemiology website: http://www.doh.state.fl.us/disease\_ctrl/epi/htopics/flu/FluLabReportGuide.pdf

<sup>\*</sup>Please note that lab event date is defined as the earliest of the following dates associated with the lab: date collected, date received by the laboratory, date reported, or date inserted.


As of 9:30 a.m. May 30, 2012 a total of 39 (58%) counties had reported their weekly level of influenza activity. During the summer months, counties have the option of continuing to report influenza activity. Please note that data reported by counties after the deadline Tuesday at 5 p.m. are recorded but may not be included in the activity map for previous weeks.

TABLE 3: Weekly County Influenza Activity for Week 21 (ending May 26, 2012) as Reported by 9:30 a.m. May 30, 2012

Activity Level	Week 20 Number of Counties	Week 21 Number of Counties	Week 21 Counties
No Report	-	29	Baker, Brevard, Citrus, Dade, Dixie, Escambia, Franklin, Gilchrist, Gulf, Hardee, Highlands, Jefferson, Lafayette, Lee, Monroe, Nassau, Okaloosa, Okeechobee, Osceola, Palm Beach, Pinellas, Putnam, Santa Rosa, Sumter, Taylor, Union, Volusia, Walton
No Activity	37	17	Calhoun, Charlotte, Columbia, Desoto, Gadsden, Glades, Hamilton, Hendry, Hernando, Holmes, Jackson, Liberty, Madison, Manatee, St. Johns, Suwannee, Washington
Mild	28	19	Alachua, Bay, Broward, Clay, Collier, Duval, Flagler, Hillsborough, Indian River, Lake, Leon, Marion, Martin, Pasco, Polk, Sarasota, Seminole, St. Lucie, Wakulla
Moderate	2	3	Bradford, Levy, Orange
Widespread	-	-	-


Map 2: Weekly County Influenza Activity for Week 21 as Reported by 9:30 a.m. May 30, 2012

A total of 39 counties report influenza activity in week 21. Most reporting counties answered no or mild activity. Three counties report moderate activity.



County influenza activity data is reported to the Bureau of Epidemiology through EpiGateway on a weekly basis by the county influenza coordinator. Specific information is requested about laboratory results, outbreak reports, and surveillance system activity. Figures 16-25 displayed below reflect a county's assessment of influenza activity within their county as a whole as well as influenza activity within specific settings. For week 21, four counties indicated that activity was decreasing, 26 indicated it was about the same as previous weeks, and seven indicated that activity was increasing.

**FIGURE 16** shows the assessment of the overall influenza activity trend in each county as reported by county health department flu coordinators for week 21 as of 9:30 a.m. May 30, 2012.



Definitions for the County Influenza Activity Trends are available at: http://www.doh.state.fl.us/disease\_ctrl/epi/CountyInfluenzaTrendGuide.html

Counties are asked to evaluate influenza activity in certain settings within their county. Each setting has a scale for activity that ranges from none or minimal activity to very high activity. What defines each of the values varies by facility type, but the example of the assessment in elementary, middle, and high schools is included below. More detailed information on the meanings of the levels for each setting can be found on the webpage also included below.

No or very minimal activity -- Scattered cases of ILI with no increase in absenteeism or disruption of school activities.

Moderate activity -- Absenteeism elevated above baseline (in range of 10 to 25%) in some but fewer than half of schools where it is known; occasional children sent home because of ILI.

High activity -- Absenteeism elevated above baseline (in range of 10 to 25%) in more than half of schools; most schools sending several or many children home each day because of ILI.

Very high activity -- Absenteeism high enough to force curtailment of some or all school activities.

FIGURE 17: Assessment of Influenza Activity in Elementary, Middle,

County influenza settings assessment guides are available at: http://www.doh.state.fl.us/disease\_ctrl/epi/FluAssessment.htm

**FIGURE 17 - FIGURE 18** show the activity levels in various facilities by county as reported by county health department flu coordinators for week 21 as of 9:30 a.m. May 30, 2012.

And High Schools

Very High

High

None or Minimal

Not Applicable

No answer

1

No answer

No moderate

1

No answer

No answer

No moderate

1

No answer

No answer

No answer

O 10 20 30 40 50 60 70

Number of Counties




FIGURE 19 - FIGURE 25 show the activity levels in various facilities by county as reported by county health department flu coordinators for week 21 as of 9:30 a.m. May 30, 2012.

