

Week 39: September 23 – 29, 2012

Summary

The Florida Department of Health (FDOH) monitors multiple surveillance systems such as the Electronic Surveillance System for the Early Notification of Community-based Epidemics (ESSENCE), the Florida Pneumonia and Influenza Mortality Surveillance System (FPIMSS), notifiable disease reports (Merlin), EpiCom, and Florida ILINet in order to track influenza activity in the state. **National:**

Since July, 2012, 307 people in 11 states have been infected with a variant strain of influenza A H3N2, also known as H3N2v. The H3N2v strain originates from influenza that circulates in pigs. Nearly all persons infected with H3N2v have reported direct or indirect contact with swine. Most cases have occurred in children; one death has been reported in an adult with underlying health conditions. **No cases of H3N2v infection have been reported in Florida.** More information can be found at: <u>http://</u>www.cdc.gov/flu/swineflu/influenza-variant-viruses-h3n2v.htm

State:

After experiencing elevated influenza-like illness (ILI) levels statewide during the summer months, ESSENCE emergency department ILI levels have increased in recent weeks. The sharpest increases in ESSENCE ILI have been seen in Regions 2, 3 and 5. One county reported moderate influenza activity in week 39. Alachua county reported an ILI outbreak in an elementary school in week 39. Five students had ILI symptoms. One student tested rapid antigen positive for influenza B; another student tested rapid antigen positive for influenza A.

In week 39, three specimens tested PCR positive for Influenza A H3 at the state public health laboratory. In recent weeks, influenza B has been the most common influenza subtype identified by the state public health laboratory. RSV activity is elevated above previous years at this time although recent increases are in keeping with seasonal trends.

TABLE 1: Summary of Florida Influenza-Like Illness (ILI) Activity for Week 39

Measure	Difference from previous week	Current week 39	Previous week 38	Page of report	
Percent of visits to ILINet providers for ILI	▲ 0.1	1.2%	1.1%	2	
Number of ILI outbreaks reported in EpiCom	▲ 1	1	0	2	
Number of counties reporting moderate or widespread influenza activity	No Change	1	1	2	
Percent of emergency department visits (from ESSENCE) due to ILI	▲ 0.5	2.7%	2.2%	3	
Proportion of laboratory specimens that were positive for influenza out of total submissions	▼ 5.5%	11.1%	16.6%	5	


NREVSS Respiratory Virus Surveillance

The National Respiratory and Enteric Virus Surveillance System (NREVSS) collects data from laboratories around the country on a weekly basis. NREVSS monitors temporal and geographic patterns of six common respiratory viruses. Fifteen facilities reported in week 39.

FIGURE 1 shows the percentage of positive tests for multiple respiratory viruses reported by NREVSS-participating laboratories in Florida.

The 6 respiratory viruses summarized in Figure 1 are:

- Respiratory Syncytial Virus (RSV)
- Parainfluenza 1-3
- Adenovirus
- Human Metapneumovirus (HMPV)
- Rhinovirus
- Influenza



HEALTH

Posted on the Bureau of Epidemiology website: <u>http://</u> www.doh.state.fl.us/floridaflu/

October 3, 2012


Produced by: Bureau of Epidemiology, Florida Department of Health (FDOH)

Contributors: Heather Rubino, MS; Colin Malone, MPH; Laura Coleman, BS; Aaron Chern, MS; Leah Eisenstein, MPH; Lea Heberlein-Larson, BS, MPH, SM(ASCP); Valerie Mock; Janet Hamilton, MPH


In this Issue:	
Summary	1
NREVSS Respiratory Virus Surveillance	1
Outpatient Influenza-like Illness Surveillance Network (ILINET)- Statewide	2
Influenza and ILI outbreaks reported in EpiCom	2
County Influenza Activity	2
ESSENCE Syndromic Surveillance Summary-Statewide	3
ESSENCE Syndromic Surveillance Summary-Regional	4
Florida Bureau of Public Health Laboratories Viral Surveillance	5

ILINET Influenza-like Illness-Statewide

ILINet is a nationwide surveillance system composed of sentinel providers. Florida has 110 providers enrolled in ILINet who submit weekly ILI and total visit counts, as well as submitting ILI specimens to the Bureau of Public Health laboratories (BPHL) for confirmatory testing.



*ILI = Influenza-like illness, fever >100°F AND sore throat and/or cough *in the absence* of another known cause. ^There is no week 53 during the 2009-2010, 2010-2011, and 2011-12 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1.



†Data presented here are counts, not proportions as included in Figure 2. This is because age group denominator data is not available through ILINet.

Influenza and ILI Outbreaks

H3N2v, FDOH is enhancing virologic

surveillance. ILI Sentinels have been

advised to submit more specimens to BPHL from children under age 18.

BPHL is able to detect presumptive

testing. No cases of variant H3N2 influenza A have been reported in

Florida. Detailed guidance

positives for both seasonal and novel influenza viruses through routine

documents have been sent to sentinel

sites and county health departments.

FIGURE 3 shows ILI visit counts

statewide by age group.

reported by ILINet sentinel providers

In week 39, ILI visit counts decreased

in the 0-64 age group. ILI visit counts

are flat in the 65+ age group.

In week 39, there was one influenza or ILI outbreak reported via EpiCom.

• Alachua County: An elementary school reported that five students had ILI. One student tested positive for influenza B by rapid antigen test. Another specimen tested positive for influenza A by rapid antigen test. None of the five students had been vaccinated.

Nine influenza or ILI outbreaks have been reported via EpiCom in summer 2012.

County Influenza Activity


As of 10:00 a.m. October 2, 2012 a total of 36 (54%) counties had reported their weekly level of influenza activity. Levy County reported moderate activity; the other 31 counties reported mild or no activity. During the summer months, counties have the option of continuing to report influenza activity. *Please note that data reported by counties after the deadline Tuesday at 5:00 p.m. are recorded but may not be included in the activity map for previous weeks.*

ESSENCE Syndromic Surveillance Summary-Statewide

Florida uses ESSENCE for syndromic surveillance, which currently collects data daily from 174 hospital emergency departments (EDs). These data are processed into 11 different syndrome categories based on the patient's chief complaint. One of the categories is ILI, which is composed of chief complaints that include the words "influenza" or "flu," or complaints that contain fever plus cough and/or sore throat.

FIGURE 4 shows ESSENCE data on ILI visits to EDs as a percentage of all ED visits.


Overall activity for ILI reported in ESSENCE is elevated compared to levels seen in previous non-pandemic seasons at this time.



*There is no week 53 for the 2007-2008, 2009-2010, and 2010-2011 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1.

FIGURE 5 shows percentage of ILI among all ED visits by age group.

Age-specific trends show that percent ILI in all age groups are increasing in week 39.



One hundred five ESSENCE participating facilities are able to provide discharge disposition data for their ED visits going back to week 40, 2010. Using this information, the percent of ED visits for ILI that result in hospitalization can be calculated. The highest percentage of admissions is in the 55+ years old age group. The low number of visits in the 55+ age group causes variability in the ILI admission percentage from week to week.

FIGURE 6 shows the percentage of ED visits for ILI that resulted in hospitalization, by age group.




FIGURE 6: Percentage of ED Visits for ILI Resulting in Hospital Admission, Hospitals Reporting Discharge Disposition (N=105), Week 40, 2010 to Week 39, 2012

ESSENCE Syndromic Surveillance Summary-Regional

page 4

FIGURE 7 - FIGURE 13 describe ED chief complaint data from ESSENCE by Regional Domestic Security Task Force (RDSTF) region.

ILI activity in ESSENCE is elevated over previous years at this time in Regions 1, 2, 3, and 5.









FIGURE 8: Percentage of Influenza-Like Illness from Emergency Department (ED) Chief Complaints, RDSTF Region 2 ESSENCE Participating Hospitals (N=2), Week 40, 2009 through October 2, 2012













FIGURE 9: Percentage of Influenza-Like Illness from Emergency Department (ED) Chief Complaints, RDSTF Region 3 ESSENCE Participating Hospitals (N=14), Week 40, 2009 through October 2, 2012






FIGURE 13: Percentage of Influenza-Like Illness from Emergency Department (ED) Chief Complaints, RDSTF Region 7 ESSENCE Participating Hospitals (N=48), Week 40, 2009 through October 2, 2012 8 Percentage of ED Visits 6 4 2 0 2 2 4 4 4 46 48 50 52 Ξ 2 Week -2009-2010* Percent -2010-2011 Percent -2011-2012 Percent -1-2012-2013 Percent

Return to Top

*There is no week 53 for the 2009-2010, 2010-2011 and 2011-2012 seasons; the week 53 data point for those seasons is an average of weeks 52 and 1. **Historical data for region 2 is only available beginning week 1, 2009.

Bureau of Public Health Laboratories Viral Surveillance

TABLE 2 shows the number of specimens tested by the BPHL, how many are influenza positive, and how many are H1N1 or other influenza subtypes.

FIGURE 14 - FIGURE 15 use BPHL viral surveillance data to track the progress of influenza infection over time. They include weekly information on how many specimens are tested by the BPHL, what proportion of those test positive for influenza, and what subtypes are found for the positive.


Small numbers of influenza specimens submitted to BPHL tested positive for influenza A H3, 2009 H1N1, and influenza B.


Influenza B has been the most common strain detected by BPHL in recent weeks.




	Current Week 39	Previous Week 38
Total Specimens Tested	27	18
Influenza Positive Specimens (% of total)	3 (11.1%)	3 (16.6%)
H1N1 Positive Specimens (% of influenza positives)	-	-
H3 Influenza A (% of influenza positives)	3 (100.0%)	-
Influenza A Unspecified (% of influenza positives)	-	2 (66.7%)
Influenza B Unspecified (% of influenza positives)	-	1 (33.3%)

FIGURE 14: Number of Influenza-Positive Specimens Tested by the Florida Bureau of Public Health Laboratories (BPHL) by Subtype by Lab Event Date* Week 1, 2011 to Week 39, 2012 as Reported in Merlin by 10:00 a.m. October 3, 2012







*Please note that lab event date is defined as the earliest of the following dates associated with the lab: date collected, date received by the laboratory, date reported, or date inserted.

For county-specific laboratory data, please refer to the Flu Lab Report in Merlin.

For instructions on how to use the Flu Lab Report, please see the Guide to Flu Lab Report on the Bureau of Epidemiology website: http://www.doh.state.fl.us/disease_ctrl/epi/htopics/flu/FluLabReportGuide.pdf