

Vaccine-Preventable Disease Surveillance

February 2018

Summary

Pertussis

- Pertussis activity increased from last month and is consistent with trends seen in previous years at this time.
- There were 28 cases and no outbreaks reported in February.
- Incidence remains highest among infants less than one year old; infants less than two months old are too young to receive vaccinations against pertussis, which is why vaccination of other age groups is so important to help prevent infection in infants.

Varicella

- Varicella activity decreased from last month and is lower than trends seen in previous years at this time.
- There were 45 cases and no outbreaks reported in February.
- Incidence remains highest among infants less than one year old, who are too young to be vaccinated
- This month, 58% of cases were not up to date on their varicella vaccinations or had unknown vaccination status.


Mumps

- Mumps activity decreased from last month but remains elevated from trends seen in previous years at this time.
- There were four cases and no outbreaks reported in February.
- Incidence was highest among children age 6-11 years old.
- This month, 75% of cases were not up-to-date on their mumps vaccinations or had an unknown vaccination status.

Table of Contents

Pertussis Surveillance	2
Varicella Surveillance.....	5
Mumps Surveillance	8
Surveillance System Summary.....	11

Monthly Activity Trends Mar. 2017-Feb. 2018



For all vaccine-preventable diseases, vaccination is the best way to prevent infection. In general, those who are at least partially vaccinated against a disease, even if they later develop that disease, have less severe outcomes than those who have never been vaccinated.


Vaccination

Map 1 shows the prevalence of children age 4-18 who have a registered religious exemption (RE) to vaccination in Florida SHOTS* as of March 7, 2018.

Statewide, the estimated prevalence of REs among children age 4-18 is 2.9% with individual counties ranging from 0.4% to 8.3%.

Unvaccinated children are at increased risk of vaccine-preventable diseases like pertussis, varicella, and mumps. Additionally, communities with a higher proportion of RE are at increased risk of vaccine preventable disease transmission.

*FL SHOTS (State Health Online Tracking System) is Florida's statewide immunization registry. All REs are required to be entered into FL SHOTS.



Posted March 7, 2018 on the Bureau of Epidemiology (BOE) website: www.floridahealth.gov/VPD

Produced by the BOE, Florida Department of Health

Contributors: Katie Kendrick, MPH; Heather Rubino, PhD; Amy Bogucki, MPH; Julia Munroe, MS; Mwedu Mtenga, MPH; Lea Heberlein-Larson, MPH; Valerie Mock, BS; Marshall Cone, MPH; Pam Colarusso, MSH; Janet Hamilton, MPH; Leah Eisenstein, MPH.



Summary

February 2018


State pertussis activity:

- **Twenty-eight confirmed and probable pertussis cases** were reported among 16 counties in February.
 - Pertussis activity increased from the previous month and remains consistent with trends observed in previous years at this time.
 - From January 1, 2018 through February 28, 2018, 48 confirmed and probable cases of pertussis were reported among 21 of Florida's 67 counties.
- Since 2014, an overall decrease in the annual number of confirmed and probable cases of pertussis reported has been observed. Pertussis is naturally cyclic in nature with peaks in disease every 3-5 years.
- **There were no outbreaks of pertussis reported in February.**
 - For most pertussis cases, exposure to other known cases is never identified, and they are not able to be linked to outbreaks.
- **In February, for every pertussis case identified, there was an average of three exposed contacts who were recommended antibiotics to prevent illness.** For those diagnosed with pertussis, antibiotics can shorten the amount of time they are contagious to others. Antibiotics can also be used to prevent illness in those who have been exposed to someone with pertussis while they are contagious.
- **Infants less than one year old had the highest incidence of pertussis.** This is consistent with national trends, which also show the highest incidence rate in infants less than one year old. Infants less than two months old are too young to receive vaccinations against pertussis, which is why vaccination of other age groups is so important to help prevent infection in infants.
- **Vaccination is the best way to prevent pertussis infections.** In February, 57% of reported cases had not received the recommended number of pertussis vaccinations for their age or had unknown vaccination status. In general, those who have received at least one pertussis vaccination have less severe outcomes than those who have never been vaccinated. In February, those who were under-vaccinated were most likely to visit the emergency department and require inpatient hospitalization.
- To learn more about pertussis, please visit <http://www.floridahealth.gov/pertussis>.

National pertussis activity:

- The number of pertussis cases has been gradually increasing since the 1980s, peaking in 2012 at levels not seen since the 1950s. Since 2012, the number of pertussis cases has started to gradually decrease.
- Pertussis incidence has remained highest among infants less than one year old and lowest among those age 20 and older since the 1990s.

Map 2 Average Pertussis Incidence Rates per 100,000 Population, November 2017 through January 2018



Pertussis surveillance goals:

- Pertussis surveillance is conducted to identify cases to limit transmission in settings with infants or others who may transmit pertussis to infants, and identify and prevent outbreaks.
- Surveillance is also conducted to identify contacts of cases and recommend appropriate prevention measures, including exclusion, antibiotic prophylaxis and immunization and to monitor the effectiveness of immunization programs and vaccines. For more information on the data sources used in Florida for pertussis surveillance, see page 11 ►

Pertussis Cases by Month Reported




Figure 1 shows the number of confirmed and probable cases of pertussis reported into Merlin, January 2018 through February 2018 and the previous five-year average.

In February, the number of reported pertussis cases increased from January but remained below the five-year average. In general, the number of reported pertussis cases tends to be highest during the summer months.


Pertussis Outbreaks

Figure 2 shows the number of confirmed and probable cases that were associated with at least one other case and the total number of confirmed and probable cases as reported into Merlin, February 2018 and the previous three-month average. Cases associated with at least one other case are shown by type of association.

In February, four (14%) cases were associated with other cases through living in the same household.

Outbreak Summary:

In February, no outbreaks of pertussis were reported. In 2017, a total of five outbreaks of pertussis were reported, all in school settings.



Pertussis Treatment and Contacts





Figure 3 shows the number of confirmed and probable cases of pertussis, as reported into Merlin, and the number of contacts who were recommended antibiotics to prevent illness, February 2018 and 2018 to date.


On average, for each case reported in February there were three people exposed to the case who were recommended antibiotics to prevent illness.



Pertussis Age-Specific Incidence Rates

Figure 4 shows the age-specific incidence rates of confirmed and probable cases of pertussis, as reported into Merlin, November 2017 through February 2018.

In February, the incidence rate was highest among infants less than one year old, which is consistent with previous months. Infants experience the greatest burden of pertussis infections, not only in number of cases but also in severity. Infants less than two months old are too young to receive vaccinations against pertussis, which is why vaccination of grandparents, parents, siblings, and other age groups is so important to help prevent infection in infants.




Vaccination History for Pertussis Cases

UTD = up-to-date

Figure 5 shows the vaccination status of pertussis cases by age group for confirmed and probable cases of pertussis, as reported into Merlin, January 2018 through February 2018 (n=48).

Two (67%) individuals age four to five months and eight (47%) individuals age 19 and older were not up-to-date on their pertussis vaccinations. The majority of individuals in all other age groups were up-to-date on their pertussis vaccinations.





Pertussis Cases in Vaccinated Individuals

UTD = up-to-date

Figure 6 shows the percent of confirmed and probable pertussis cases who were up-to-date on their pertussis vaccinations, as reported into Merlin, January 2018 through February 2018 and the previous five-year average. **Figure 7** shows the percent of these cases who were under vaccinated during the same time periods.

Although individuals who have been vaccinated can still get pertussis, vaccination remains the best way to prevent pertussis and severe complications.



Pertussis Outcomes

UTD = up-to-date, ED = emergency department




Figure 8 shows the percent of confirmed and probable cases of pertussis with select outcomes by vaccination status, as reported into Merlin, February 2018 and the previous three-month average.

In February, cases who were under-vaccinated were more likely to visit the emergency department and require inpatient hospitalization.

In general, older individuals are more likely to experience paroxysmal cough while younger individuals are more likely to experience posttussive vomiting and whoop. Primarily infants less than one year old experience apnea.

Summary

February 2018

State varicella activity:

• **Forty-five confirmed and probable varicella cases were reported among 27 counties in February.**

- The number of reported varicella cases decreased slightly from January and is lower than trends seen in previous years at this time. In previous years, varicella activity was highest during the late winter and spring; trends for 2018 will continue to be monitored closely.
- From January 1, 2018 through February 28, 2018, 92 cases of varicella were reported among 34 of Florida’s 67 counties.

• A decreasing trend in the number of confirmed and probable cases of varicella reported annually in Florida was observed from 2008-2014. The number of cases reported annually remained elevated in 2015 and 2016 and started to decrease again in 2017. So far in 2018, the number of varicella cases is lower than that observed in 2017 at this time.

• **No outbreaks of varicella were reported in February.**

• **In February, children age less than one year old had the highest incidence of varicella.** This is consistent with what was observed for the majority of months in 2017.


• **Vaccination is the best way to prevent varicella infection.** In February, 58% of cases were not up to date on their varicella vaccinations or had unknown vaccination status. In general, those who received at least one dose of varicella vaccination, even if they later develop disease, have less severe outcomes than those who have never been vaccinated. In February, infants too young for vaccination were most likely to visit the emergency department.

• To learn more about varicella, please visit <http://www.floridahealth.gov/varicella>.

National varicella activity:

- Varicella incidence decreased significantly following the vaccine becoming available in 1995 and has continued to decrease since 2006 when recommendations changed from one to two doses of varicella vaccine.
 - From 2006 –2015 all age groups saw a significant decrease in incidence with the largest decline in children age 5-9 years and age 10-14 years.
- Although varicella is not reportable in all states and therefore not all states report varicella cases to the CDC, based on available data the number of varicella cases nationally has steadily decreased each year from 2012-2015.

Map 3 Average Varicella Incidence Rates per 100,000 Population, November 2017 through January 2018



Surveillance goals:

- Varicella surveillance is conducted to identify and control outbreaks and monitor trends and severe outcomes.
- Surveillance is also conducted to monitor effectiveness of immunization programs and vaccines. For more information on the data sources used in Florida for varicella surveillance, see page 11 ►

Varicella Cases by Month Reported




Figure 9 shows the number of confirmed and probable cases of varicella reported into Merlin, January 2018 through February 2018 and the previous five-year average.

In February, the number of reported varicella cases decreased slightly from January and remained below the previous five-year average. Thus far in 2018, varicella activity does not appear to be following seasonal trends seen in previous years at this time.


Varicella Outbreaks

Figure 10 shows the number of confirmed and probable cases that were associated with at least one other case and the total number of confirmed and probable cases as reported into Merlin, February 2018 and the previous three-month average. Cases associated with at least one other case are shown by type of association.

In February, six (13%) cases were associated with other cases through living in the same household.

Outbreak Summary:

No outbreaks of varicella were reported in February. Thus far in 2018, two varicella outbreaks have been reported in correctional facilities.



Varicella Age-Specific Incidence Rates




Figure 11 shows the age-specific incidence rates of confirmed and probable cases of varicella, as reported into Merlin, November 2017 through February 2018.


In February, the incidence rate was highest among infants less than one year old. This is consistent with trends seen in 2017. Infants less than one year old are too young to receive varicella vaccination, which is why vaccination of other age groups is so important to help prevent infection in infants.

Vaccination History for Varicella Cases

UTD = up-to-date

Figure 12 shows the vaccination status of varicella cases by age group for confirmed and probable cases of varicella, as reported into Merlin, January 2018 through February 2018 (n=92).

Varicella vaccinations are recommended at 12-15 months of age and 4-6 years of age. Twenty-two (46%) individuals age 19 and older were not up-to-date on their varicella vaccinations, while the majority of those age 1-11 years were up-to-date on their vaccinations.





Varicella Cases in Vaccinated Individuals

UTD = up-to-date

Figure 13 shows the percent of confirmed and probable varicella cases who were up to date on their varicella vaccinations, as reported into Merlin, January 2018 through February 2018 and the previous five-year average. **Figure 14** shows the percent of these cases who were under vaccinated during the same time periods.

Although individuals who have been vaccinated can still get varicella, vaccination remains the best way to prevent varicella and severe complications.




Varicella Outcomes

UTD = up-to-date, ED = emergency department

Figure 15 shows the percent of confirmed and probable cases of varicella with select outcomes by vaccination status, as reported into Merlin, February 2018 and the previous three-month average.

In general, cases who were UTD on their vaccinations were less likely to experience fever and vesicle lesions. Cases too young to be vaccinated or not yet vaccinated were more likely to experience papule lesions.

In February, individuals too young for vaccination were most likely to visit the emergency department. Few varicella cases require inpatient hospitalization. In general, those who received at least one dose of varicella vaccination, even if they later develop disease, have less severe outcomes than those who have never been vaccinated.



Summary

February 2018

State mumps activity:

- **Two confirmed and two probable mumps cases were reported among three counties in February.**
 - Mumps cases have remained elevated since April 2017 with a peak of 20 cases reported in August 2017.
 - From January 1, 2018 through February 28, 2018, nine confirmed and five probable cases of mumps were reported among five of Florida's 67 counties.
- In Florida, the number of reported mumps cases has remained relatively low over the past five years but has steadily increased since 2015 (10 cases), with a large spike in 2017 (70 cases). The last time the number of reported cases reached 2017 levels was in the 1990s.
- **No outbreaks of mumps were reported in February.**
 - In 2017, the majority of mumps cases were associated with outbreaks or household clusters.
 - While mumps outbreaks can occur in highly-vaccinated communities, high vaccination coverage limits the size, duration, and spread of outbreaks.
- **In February, the highest incidence of mumps was in children age 6-11 years old.**
- **Vaccination is the best way to prevent mumps infections.** In February, 75% of cases were not up-to-date on their mumps vaccinations or had an unknown vaccination status.
- In February, one (25%) unvaccinated individual was hospitalized for their illness. In general, those who have received at least one mumps vaccination even if they later develop disease have less severe outcomes than those who have never been vaccinated.
- To learn more about mumps, please visit <http://www.floridahealth.gov/mumps>.

National mumps activity:

- Since 1989 when the two dose vaccination program was introduced, the number of mumps cases has fluctuated from a few hundred to a few thousand per year. Some years had higher numbers of cases than others mainly because of several large outbreaks in close-contact settings.
- In 2016, there were over 6,000 cases of mumps reported, and in 2017 there were over 5,600 cases reported. Since 2013, the 18-22 year age group has had the highest incidence of mumps, largely driven by outbreaks. About half of the outbreaks reported since 2016 have been associated with colleges and universities, primarily affecting young adults.

Map 4 Cumulative Mumps Incidence Rates per 100,000 Population, February 2018





Figure 16 shows the number of confirmed and probable cases of mumps reported into Merlin, 2013 through February 2018.



Surveillance goals:

- Mumps surveillance is conducted to identify and control outbreaks and monitor trends and severe outcomes.
- Surveillance is also conducted to monitor effectiveness of immunization programs and vaccines. For more information on the data sources used in Florida for mumps surveillance, see page 11 ►

Mumps Cases by Month Reported




Figure 17 shows the number of confirmed and probable cases of mumps reported into Merlin, January 2018 through February 2018 and the previous five-year average.

In February, the number of reported mumps cases decreased from that in January, but remained above the previous five-year average. Cases have been elevated since the summer months of 2017, peaking in August and December when several cases associated with outbreaks and household clusters were reported.


Mumps Outbreaks

Figure 18 shows the number of confirmed and probable cases that were associated with at least one other case and the total number of confirmed and probable cases as reported into Merlin, February 2018 and the previous three-month average. Cases associated with at least one other case are shown by type of association.

In February, all reported mumps cases were sporadic and not associated with any other reported cases.

Outbreak Summary:

No mumps outbreaks were reported in February. There have been no mumps outbreaks reported thus far in 2018.



Mumps Age-Specific Incidence Rates




Figure 19 shows the age-specific incidence rates of confirmed and probable cases of mumps, as reported into Merlin, November 2017 through February 2018.


In February, the incidence rate was highest among children age 6-11 years old. In recent months, the majority of cases have been in children age 12-18 years and adults age 19 years and older.

Vaccination History for Mumps Cases

UTD = up-to-date

Figure 20 shows the vaccination status of mumps cases by age group for confirmed and probable cases of mumps, as reported into Merlin, January 2018 through February 2018 (n=14).

Mumps vaccinations are recommended at 12-15 months of age and 4-6 years of age. Two (25%) individuals age 19 years and older were not up-to-date on their mumps vaccinations, while the majority of individuals age 6-18 years old were up-to-date on their vaccinations.





Mumps Cases in Vaccinated Individuals

UTD = up-to-date

Figure 21 shows the percent of confirmed and probable mumps cases who were up to date on their mumps vaccinations, as reported into Merlin, January 2018 through February 2018 and the previous five-year average. **Figure 22** shows the percent of these cases who were under vaccinated during the same time periods.

Although individuals who have been vaccinated can still get mumps, vaccination remains the best way to prevent mumps and severe complications.



Mumps Outcomes


UTD = up-to-date, ED = emergency department

Figure 23 shows the percent of confirmed and probable cases of mumps with select outcomes by vaccination status, as reported into Merlin, February 2018 and the previous three-month average.

In February, two (50%) cases visited the emergency department, and one (25%) unvaccinated individual was hospitalized.

Orchitis (testicular inflammation) is the most common complication from mumps in males. From January 2018 through February 2018, two (18%) cases reported orchitis; one was never vaccinated and one had an unknown vaccination status.

In general, those who received at least one dose of mumps vaccination, even if they later develop disease, have less severe outcomes than those who have never been vaccinated.



Case Data

- Pertussis, varicella, and mumps are reportable diseases in Florida. Case information is documented by county health department (CHD) epidemiologists in Merlin, Florida's reportable disease surveillance system.
- CHD epidemiologists also report outbreaks of pertussis, varicella, and mumps into Merlin. Outbreaks are defined as two or more cases associated with a specific setting outside of the home. Two or more cases among members of the same household are considered household-associated cases.
- Current case information is preliminary and may change as more data are received. The most recent data available are displayed in this report.
- For more information about reportable diseases, please visit www.Floridahealth.gov/diseasereporting.

Population Data

- Population data used to calculate incidence rates are from FLHealthCHARTS (Community Health Assessment Resource Tool Set).
- For more information about FLHealthCHARTS, please visit www.flhealthcharts.com.

Vaccination Data

- Vaccination data from cases are from Merlin, as identified by CHD epidemiologists.
- Vaccination status is determined using the Advisory Committee on Immunization Practices Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger, 2018.
- Cases are considered up-to-date if they have received the recommended number of doses of vaccine for a particular disease for their age at the time of their illness onset. Cases are considered under vaccinated if they have received at least one but not all doses of vaccine recommended for a particular disease for their age at the time of their illness onset.
- For more information about immunization schedules, please visit <https://www.cdc.gov/vaccines/schedules/index.html>.