

Hepatitis A Surveillance

2018-To-Date Key Points

 5,024 cases

 23% cases linked to other cases



 Since 2018, 30-39 years olds had highest incidence rate

 24% co-infected with hepatitis B or C




The number of reported hepatitis A cases in February decreased from the previous month and was below the previous 5-year average. Since January 1, 2018, 98% of cases have likely been acquired in Florida.

In February, 26 hepatitis A cases were reported in **10 counties**, outlined in black in the map below. In the past three months, North Florida had the highest average incidence rates.



Since January 1, 2021, 62 hepatitis A cases were reported.

Due to the national hepatitis A outbreak, cases increased dramatically during 2018 and 2019 in Florida. In February 2021, there was a 79% decrease in overall cases when compared to cases in February 2019.



*The white bars indicate the total number of cases as of February for each year



98%
never vaccinated

The best way to prevent hepatitis A infection is through vaccination. Since January 1, 2018, 98% of people with hepatitis A had never received a documented dose of hepatitis A vaccine. In February 2021, 100% of infected people had not received the vaccine. Since 2006, hepatitis A vaccine has been recommended for all children at age 1 year. Hepatitis A vaccine is also recommended for certain high-risk groups of adults including injection and non-injection drug use, persons experiencing homelessness, and men who have sex with men. To learn more about the hepatitis A vaccine, talk to your doctor or visit: www.CDC.gov/Vaccines/HCP/VIS/VIS-Statements/Hep-A.html.


The COVID-19 pandemic is affecting health care seeking behavior, which may be impacting the diagnosis and reporting of hepatitis A cases that are shown in this report. For more information on the COVID-19 pandemic in Florida, please visit FloridaHealthCOVID-19.gov.



Hepatitis A Surveillance




Since 2018, **1,147 (23%)** of 5,024 hepatitis A cases were **epidemiologically (epi) linked to other cases**. In February 2021, 19% of cases were epi-linked to other cases.




In February 2021, all epi-links were **household contacts**.

There were an **average of 2 contacts per case**. Contacts are those who were exposed to the virus and recommended prophylaxis for illness prevention.



Since 2018, **incidence rates have increased among all age groups** with cases reported primarily among **men (64%)** and persons who identify as **non-Hispanic white (82%)**. As of February 2021, **30-39 years old had the highest incidence rate at 0.7 cases per 100,000 population**.



Since 2018, **91 cases (2%)** were co-infected with chronic hepatitis B, **1003 cases (20%)** were co-infected with chronic hepatitis C, and **106 cases (2%)** were co-infected with both chronic hepatitis B and C. In February 2021, 5 cases (19%) were co-infected with chronic hepatitis B or C. Co-infection with more than 1 type of viral hepatitis can lead to more severe liver disease and increase the risk of **developing** liver cancer.



National activity

Hepatitis A rates have decreased by more than 95% since the first vaccine became available in 1995. However, since March of 2017, the Centers for Disease Control and Prevention has been monitoring outbreaks in 35 states among persons who use drugs and persons who are experiencing homelessness. More information about these outbreaks can be found here: www.cdc.gov/hepatitis/outbreaks/2017April-HepatitisA.htm

Hepatitis A surveillance goals

- Identify and control outbreaks and monitor trends
- Identify and mitigate common sources
- Monitor effectiveness of immunization programs and vaccines

To learn more about hepatitis A, please visit FloridaHealth.gov/HepA. For more information on the data sources used in Florida for hepatitis A surveillance, see the last page of this report.


Hepatitis A Surveillance

Statewide Response to the Increase in Hepatitis A Cases

Several Florida counties have experienced ongoing local transmission of hepatitis A since 2017. Since 2018, 98% of Florida's cases (n=5,024) have likely been acquired in Florida. These cases share several common risk factors including drug use (both injection or non-injection drugs), identifying as men who have sex with men, or recently experiencing homelessness. **Individuals with any of these risk factors should receive the hepatitis A vaccine, and health care providers are encouraged to actively offer the hepatitis A vaccine to individuals at risk. Vaccination is the best way to prevent hepatitis A infection.**

For additional information, please see the declaration of public health emergency issued by the State Surgeon General in August 2019, available at: [FloridaHealth.gov/_documents/newsroom/press-releases/2019/08/phe-hav-filed-08-01-2019.pdf](https://www.floridahealth.gov/_documents/newsroom/press-releases/2019/08/phe-hav-filed-08-01-2019.pdf).

Since 2018, over half (61%) of the 5,024 cases acquired in Florida reported at least one of the risk factors below, while 39% reported no or unknown risk factors. The most commonly identified risk factor was **drug use**, reported by 2,753 cases (56%). Non-injection (34%) and injection (36%) were both common forms of drug use reported, followed by homelessness (18%).



Hepatitis A infections can be severe, leading to inpatient hospitalization and sometimes death. Since January 1, 2018, 3,417 (69%) cases acquired in Florida have been hospitalized due to hepatitis A infection with 76 deaths identified as hepatitis A associated.

69% hospitalized
76 deaths

The Florida Department of Health is actively working to vaccinate those most at risk for hepatitis A infection. In February 2021, 2,070 doses were administered. **The number of first doses of hepatitis A vaccine administered by both private providers and county health departments to adults age 18 years and older, as recorded in Florida SHOTS, decreased and was below the previous 5-year-average.** This may be due to changes in vaccine administration during the COVID-19 pandemic. **Vaccination is the best way to prevent hepatitis A infection.**

