Hepatitis A Surveillance

May Key Points



45 cases



13% of cases linked to other cases




30-39 year olds had the highest incidence




93% of cases not upto-date or unknown vaccination status




The number of reported hepatitis A cases in May decreased from the previous month and was below the previous 5-year average.



In May 2022, 45 hepatitis A cases were reported in 16 **counties**, outlined in black in the map below. From March 2022 through May 2022, the average county rates were highest in Central Florida and Northwest Florida.



In 2022, 215 hepatitis A cases* were reported.



^{*}The white bars indicate the total number of cases as of May for each year



not vaccinated


The best way to prevent hepatitis A infection is through vaccination. In May 2022, 93% of infected people had not received the vaccine or had unknown vaccination status. Since 2006, hepatitis A vaccine has been recommended for all children at age 1 year. Hepatitis A vaccine is also recommended for certain adult high-risk groups, including persons using injection and non-injection drugs, persons experiencing homelessness, and men who have sex with men. To learn more about the hepatitis A vaccine, talk to your doctor or visit: www.CDC.gov/Vaccines/HCP/VIS/VIS-Statements/Hep-A.html.




Hepatitis A Surveillance



In May 2022, **6 cases** were **epidemiologically (epi) linked to another case**. From January 2022—May 2022, **26 cases** were epi-linked to other cases.



In May 2022, there was an average of **6 contacts to reported cases**. Contacts are those who were exposed to the virus and recommended prophylaxis for illness prevention.



2022



In May 2022, **30-39 year olds have the highest incidence rate at 2.84 cases per 100,000 population.** Similarly in 2021, 30-39 year olds also had the highest incidence rate at 1.48 cases per 100,000 population.





In May 2022, 1 cases was co-infected with chronic hepatitis C and no cases were co-infected with chronic hepatitis B. In 2021, the most common coinfection was with chronic hepatitis C with 15% of reported cases being coinfected. Co-infection with more than one type of viral hepatitis can lead to more severe liver disease and increase the risk of developing liver cancer.

Chronic hepatitis B Chronic hepatitis C Chronic hepatitis B and C No co-infection

1% 15% 1% 83%



National activity

Hepatitis A rates have decreased by more than 95% since the first vaccine became available in 1995. However, since outbreaks were first identified in 2016, the Centers for Disease Control and Prevention has been monitoring outbreaks in 37 states. More information about these outbreaks can be found here: www.cdc.gov/hepatitis/outbreaks/2017March-HepatitisA.htm

Hepatitis A surveillance goals

- · Identify cases to limit transmission
- Identify and prevent outbreaks
- Monitor effectiveness of immunization programs and vaccines


To learn more about hepatitis A, please visit FloridaHealth.gov/HepA. For more information on the data sources used in Florida for hepatitis A surveillance, see the last page of this report.

Hepatitis A Surveillance

Vaccination is the best way to prevent hepatitis A infection. Health care providers are encouraged to actively offer the hepatitis A vaccine to individuals at risk including men who have sex with men.



In 2022, 57% of the 205 cases* in Florida reported at least one of the risk factors below, while 43% reported no or unknown risk factors. The most commonly identified risk factor was men who have sex with men, reported by 94 cases (64%). The next most common risk factor was any drug use (33%) reported in 49 cases. The most common form of drug use was non-injection drug (28%) reported in 42 cases. Injection drug use (6%) was reported in 9 cases. Recent homelessness was reported in (7%) of reported cases. In 2022, there has been an increase in reported cases among men who have sex with men when compared to 2021.






Hepatitis A infections can be severe, leading to inpatient hospitalization and sometimes death. In 2022, 141 cases (69%) reported in Florida* have been hospitalized due to hepatitis A infection. No deaths have been identified as hepatitis A associated in 2022.



The Florida Department of Health is actively working to vaccinate those most at risk for hepatitis A infection. In May 2022, 3,955 doses were administered. The number of first doses of hepatitis A vaccine administered by both private providers and county health departments to adults age 18 years and older, as recorded in Florida SHOTS, increased and was below the previous 5-year-average. Vaccination is the best way to prevent hepatitis A infection.



^{*} The 10 cases with illness prior to 2021 are excluded from this analysis.