Hepatitis A Surveillance

April Key Points



13 cases

Jan



23% of cases linked to other cases



30-39 year olds had the highest incidence



31% of cases were not up-to-date and 46% of cases had unknown vaccination status.




The number of reported hepatitis A cases in April decreased from the previous month and was below the previous 5-year average.




Jul

In April 2023, **13 hepatitis A cases** were reported in **8 counties**, outlined in black in the map below. From February 2023 through April 2023, the average county rates were lowest throughout northwest Florida.



In 2023, 54 hepatitis A cases were reported.

May Jun



Dec

*The white bars indicate the total number of cases as of April for each year



not vaccinated

The best way to prevent hepatitis A infection is through vaccination. In April 2023, 77% of reported cases had not received the vaccine or had unknown vaccination status. Of the 76%, 31% of cases were not-up to date on hepatitis A vaccinations and 46% of cases had unknown hepatitis A vaccination status. Since 2006, hepatitis A vaccine has been recommended for all children at age 1 year. Hepatitis A vaccine is also recommended for certain adult high-risk groups, including persons using injection and non-injection drugs, persons experiencing homelessness, and men who have sex with men. To learn more about the hepatitis A vaccine, talk to your doctor or visit: www.CDC.gov/Vaccines/HCP/VIS/VIS-Statements/Hep-A.html.



Hepatitis A Surveillance



In April 2023, 3 cases were epidemiologically (epi) linked to another case.




In April 2023, there was an average of **1 contact to reported cases**. Contacts are those who were exposed to the virus and recommended prophylaxis for illness prevention.





In 2023, 30-39 year olds have the highest incidence rate at 0.49 cases per 100,000 population.





In April 2023, no cases were co-infected with chronic hepatitis C and no cases were co-infected with chronic hepatitis B. In 2022, most cases (97%) had no co-infection. **Co-infection with more than one type of viral hepatitis can lead to more severe liver disease and increase the risk of developing liver cancer**.

Chronic hepatitis B Chronic hepatitis C Chronic hepatitis B and C No co-infection

1% 2% 97%



National activity

Hepatitis A rates have decreased by more than 95% since the first vaccine became available in 1995. However, since outbreaks were first identified in 2016, the Centers for Disease Control and Prevention has been monitoring outbreaks in 37 states. More information about these outbreaks can be found here: www.cdc.gov/hepatitis/outbreaks/2017March-HepatitisA.htm

Hepatitis A surveillance goals

- · Identify cases to limit transmission
- Identify and prevent outbreaks
- Monitor effectiveness of immunization programs and vaccines

To learn more about hepatitis A, please visit <u>FloridaHealth.gov/diseases-and-conditions/vaccine-preventable-disease/hepatitis-a</u>. For more information on the data sources used in Florida for hepatitis A surveillance, see the last page of this report.


Hepatitis A Surveillance

Vaccination is the best way to prevent hepatitis A infection. Health care providers are encouraged to actively offer the hepatitis A vaccine to individuals at risk including men who have sex with men.

For more information about hepatitis A vaccination in Florida visit: https://www.floridahealth.gov/diseases-and-conditions/hepatitis-vaccination-testing-program.html



In 2023, 41% of the 54 cases in Florida reported at least one of the risk factors below, while 59% reported no or unknown risk factors. The most commonly identified risk factor was any drug use, reported by 17 cases (31%). The next most common risk factor was non-injection drug use (13%) reported in 7 cases followed by injection drug use (11%) reported in 6 cases. Men who have sex with men was reported in 15% of reported cases, followed by recent homelessness reported in 6% of cases.





Hepatitis A infections can be severe, leading to inpatient hospitalization and sometimes death. In 2023, 22 cases (54%) reported in Florida have been hospitalized due to hepatitis A infection. No deaths have been identified as hepatitis A associated in 2023.



The Florida Department of Health is actively working to vaccinate those most at risk for hepatitis A infection. In April 2023, 4,716 doses were administered. The number of first doses of hepatitis A vaccine administered by both private providers and county health departments to adults age 18 years and older, as recorded in Florida SHOTS, decreased and was below the previous 5-year-average. Vaccination is the best way to prevent hepatitis A infection.

