
Pertussis Surveillance

January Key Points











<1 year olds had the highest 6 month average incidence



In January 2024, 13 pertussis cases were reported in 9 counties. The number of pertussis cases reported in January decreased from the previous month and was below the previous 5-year average.



In 2024, 13 pertussis cases were reported. There was a **65% increase** in the number of pertussis cases reported between August 2023—January 2024 (n=56) compared to August 2022—January 2023 (n=34).




^{*}The white bars indicate the total number of cases as of January for each year



In January 2024, five pertussis cases were household-associated. No pertussis cases were outbreak-associated. In the past 6 months, there was an average of 2 household-associated cases, an average of 0 outbreak-associated cases and an average of 10 total cases. From August 2022 to January 2023, there was an average of 1 household-associated cases, an average of 0 outbreak-associated cases and an average of 6 total cases. For most pertussis cases, exposure to other known cases is not identified and are not able to be linked to outbreaks.

Household-associated Outbreak-associated Total cases





Pertussis Surveillance



There was an average of 2 contacts per case between August 2022 and January 2022 and between August 2023 and January 2024. Contacts are classified as people whom antibiotics were recommended to prevent illness. Antibiotics can shorten the amount of time cases are contagious and can also be used to prevent illness in those exposed. Understanding pertussis transmission is a key factor in decreasing pertussis infections. In Florida, transmission setting is not routinely identified for non-outbreak cases, resulting in 69% of cases reporting unknown setting in the past six months.

August 2022 to January 2023


August 2023 to January 2024



The average incidence rate was highest among <1 year olds at 0.352 cases per 100,000 population between August 2023 and January 2024. Infants experience the greatest burden of pertussis infections, not only in number of cases but also in severity. Infants <2 months old are too young to receive vaccinations against pertussis, which is why vaccination of parents, siblings, grandparents, and other age groups is important in infection prevention among infants.

August 2022 to January 2023


August 2023 to January 2024



Pertussis Surveillance



In 2024, over half of cases reported were not up-to-date on their pertussis vaccinations. In general, those who have received at least one pertussis vaccination have less severe outcomes than those who have never been vaccinated. If a person was born before January 1st, 1982, the current pertussis immunization recommendation would not have been implemented when they were receiving their childhood immunizations. Based on the case's age, **2 cases** would not have been vaccinated under the current childhood immunization recommendations.





National activity

The number of pertussis cases gradually increased since the 1980s, peaking in 2012 at levels not seen since the 1950s. Since 2012, the number of pertussis cases started gradually decreasing. Pertussis incidence has remained highest among infants <1 year old and lowest among adults ≥20 years old since the 1990s.

Pertussis surveillance goals

- Identify cases to limit transmission in settings with infants or others who may transmit pertussis to infants
- Identify and prevent outbreaks
- Identify transmission settings in non-outbreak cases to prevent the spread of sporadic cases
- Identify contacts of cases and recommend appropriate prevention measures, including exclusion, antibiotic prophylaxis, and immunization
- Monitor the effectiveness of immunization programs and vaccines

To learn more about pertussis, please visit FloridaHealth.gov/Pertussis. For more information on the data sources used in Florida for pertussis surveillance, see the last page of this report.