

Test Menu

Test Name (s) N/A Analyte(s) mumps Test Code 1660, 1756 (MMR Index Value Panel) Lab Iocation Jacksonville location Department Virology Prior Authorization None Required Forms Test Requisition Form, DH1847. Specimen Sources Freshly drawn serum. Supplemental Information-Special Specimen Preparation Minimum Volume 3-5 mL of blood Storage Conditions Storage Conditions Storage Conditions Collection Media Vacuations and Handling Specimen Labeling Specimen Preparation Packaging and Shipping Instructions and Handling Packaging and Shipping Instructions and Handling Serious	TOPIC	DESCRIPTION
Other Name (s) N/A Analyte(s) mumps Test Code 1660, 1756 (MMR Index Value Panel) Lab location Jacksonville location Department Virology Prior Authorization None Required Forms Test Requisition Form, DH1847. Specimen Sources Freshly drawn serum. Supplemental Information- Special Specimen Preparation N/A Minimum Volume 3-5 mL of blood Storage Conditions Store at room temperature ≤ 8 hours. May be stored between 2-8°C for a maximum of seven days, including the time for shipping and lab receipt Collection Media Vacutainer (red storeper) or serum separator tube Specimen Labeling -Specimen must be labeled with at least two unique patient identifiers, Ex: Name and DOBThe collection date and time if submitting multiple specimensInformation on the specimen must match the requisition. Packaging and Shipping Instructions and Handling We recommend sending within four days of collection. Specimens must be shipped between (2-8°C) or frozen (s-20°C) on dry ice. Separate multiple specimens into different bags (preferred). Test Methodology Serology (i.e., ELISA) Turnaround Time 7 - 14 days Result Indicator Positive, Negative or Equivocal <th></th> <th></th>		
Analyte(s) Mumps Test Code 1660, 1756 (MMR Index Value Panel) Lab location Jacksonville location Department Virology Prior Authorization None Required Forms Test Requisition Form, DH1847. Specimen Sources Freshly drawn serum. Supplemental Information-Special Specimen Preparation N/A Preparation N/A Preparation Storage Conditions Storage tonditions Store at room temperature ≤ 8 hours. May be stored between 2-8°C for a maximum of seven days, including the time for shipping and lab receipt Vacutainer (red stopper) or serum separator tube Specimen Labeling -5pecimen must be labeled with at least two unique patient identifiers, Ex: Name and DOBThe collection date and time if submitting multiple specimensInformation on the specimen must match the requisition. Packaging and Shipping Instructions and Handling We recommend sending within four days of collection. Specimens must be shipped between (2-8°C) or frozen (s-20°C) on dry ice. Separate multiple specimens into different bags (preferred). Test Methodology Serology (i.e., ELISA) Turnaround Time 7 - 14 days Result Indicator Positive, Negative or Equivocal Unsatisfactory Specimen Specimen serceived at 2-8°C or J days after collection. For storage >7 days, specimen must be frozen at s-20°C and received frozen on dry ice. Specimen must be frozen at s-20°C and received frozen on individence in Do nample and paperwork, improper collection, storage or transport of specimen, on testerally contaminated specimen, disparity between iD on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history, If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations A panel of Measles, Mumps, and Rubella IgG can be ordered under 1756, MMR Index Value.	Other Name (s)	
1660, 1756 (MMR Index Value Panel) Jacksonville location Jacksonville location Jacksonville location Jacksonville location Jacksonville location Jacksonville location Virology		
Department Virology		'
Department		
Prior Authorization Required Forms Test Requisition Form, DH1847. Specimen Sources Freshly drawn serum. N/A Nepplemental Information-Special Specimen Preparation Minimum Volume 3-5 mL of blood Storage Conditions Store at room temperature ≤ 8 hours. May be stored between 2-8°C for a maximum of seven days, including the time for shipping and lab receipt Collection Media Vacutainer (red stopper) or serum separator tube Specimen Labeling -Specimen must be labeled with at least two unique patient identifiers, Ex: Name and DOBThe collection date and time if submitting multiple specimensInformation on the specimen must match the requisition. Packaging and Shipping Instructions and Handling We recommend sending within four days of collection. Specimens must be shipped between (2-8°C) or frozen (s-20°C) on dry ice. Separate multiple specimens into different bags (preferred). Test Methodology Serology (i.e., ELISA) Turnaround Time 7-14 days Result Indicator Positive, Negative or Equivocal Unsatisfactory Specimen Specimens received at 2-8°C > 7 days after collection. For storage > 7 days, specimen must be frozen at s-20°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history, if required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations If testing a particular sample occurs early during primary infection, no detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes Additional Inform		
Test Requisition Form, DH1847. Specimen Sources Freshly drawn serum.	•	
Specimen Sources Freshly drawn serum. N/A		
Supplemental Information - Special Specimen Preparation N/A Minimum Volume 3-5 mL of blood Storage Conditions Store at room temperature ≤ 8 hours. May be stored between 2-8°C for a maximum of seven days, including the time for shipping and lab receipt Collection Media Vacutainer (red stopper) or serum separator tube Specimen Labeling -Specimen must be labeled with at least two unique patient identifiers, Ex: Name and DOB.	Specimen Sources	
Storage Conditions Store at room temperature ≤ 8 hours. May be stored between 2-8°C for a maximum of seven days, including the time for shipping and lab receipt Collection Media Vacutainer (red stopper) or serum separator tube Specimen Labeling -Specimen must be labeled with at least two unique patient identifiers, Ex: Name and DOB.		N/A
maximum of seven days, including the time for shipping and lab receipt Collection Media Vacutainer (red stopper) or serum separator tube	Minimum Volume	3-5 mL of blood
Specimen Labeling		maximum of seven days, including the time for shipping and lab receipt
Identifiers, Ex: Name and DOB. -The collection date and time if submitting multiple specimens. -Information on the specimen must match the requisition. Packaging and Shipping Instructions and Handling We recommend sending within four days of collection. Specimens must be shipped between (2-8°C) or frozen (≤-20°C) on dry ice. Separate multiple specimens into different bags (preferred). Test Methodology Serology (i.e., ELISA) Turnaround Time 7 - 14 days Result Indicator Positive, Negative or Equivocal Unsatisfactory Specimen Specimens received at 2-8°C >7 days after collection. For storage >7 days, specimen must be frozen at ≤-20°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history. If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations If testing a particular sample occurs early during primary infection, no detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes A panel of Measles, Mumps, and Rubella IgG can be ordered under 17	Collection Media	Vacutainer (red stopper) or serum separator tube
be shipped between (2-8°C) or frozen (s-20°C) on dry ice. Separate multiple specimens into different bags (preferred). Test Methodology Serology (i.e., ELISA) Turnaround Time 7 − 14 days Result Indicator Positive, Negative or Equivocal Unsatisfactory Specimen Specimens received at 2-8°C > 7 days after collection. For storage > 7 days, specimen must be frozen at ≤-20°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history. If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations If testing a particular sample occurs early during primary infection, no detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes A panel of Measles, Mumps, and Rubella IgG can be ordered under 1756, MMR Index Value. Reference Range Positive, Negative or Equivocal	Specimen Labeling	identifiers, Ex: Name and DOBThe collection date and time if submitting multiple specimens.
multiple specimens into different bags (preferred). Test Methodology Serology (i.e., ELISA) Turnaround Time 7 − 14 days Result Indicator Positive, Negative or Equivocal Unsatisfactory Specimen Specimen must be frozen at ≤-2°C >7 days after collection. For storage >7 days, specimen must be frozen at ≤-2°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history. If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations If testing a particular sample occurs early during primary infection, no detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes Additional Information & Notes Positive, Negative or Equivocal N/A	Packaging and Shipping Instructions and Handling	We recommend sending within four days of collection. Specimens must
Turnaround Time 7 − 14 days Result Indicator Positive, Negative or Equivocal Unsatisfactory Specimen Specimens received at 2-8°C > 7 days after collection. For storage > 7 days, specimen must be frozen at ≤-20°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history. If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations If testing a particular sample occurs early during primary infection, no detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes A panel of Measles, Mumps, and Rubella IgG can be ordered under 1756, MMR Index Value. Reference Range Positive, Negative or Equivocal Reference Lab N/A		
Result Indicator Positive, Negative or Equivocal Unsatisfactory Specimen Specimens received at 2-8°C >7 days after collection. For storage >7 days, specimen must be frozen at ≤-20°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history. If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations If testing a particular sample occurs early during primary infection, no detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes A panel of Measles, Mumps, and Rubella IgG can be ordered under 1756, MMR Index Value. Reference Range Positive, Negative or Equivocal Reference Lab N/A	Test Methodology	Serology (i.e., ELISA)
Unsatisfactory Specimen Specimens received at 2-8°C > 7 days after collection. For storage > 7 days, specimen must be frozen at ≤-20°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history. If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations If testing a particular sample occurs early during primary infection, no detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes A panel of Measles, Mumps, and Rubella IgG can be ordered under 1756, MMR Index Value. Reference Range Positive, Negative or Equivocal Reference Lab N/A	Turnaround Time	7 – 14 days
specimen must be frozen at ≤-20°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history. If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider, broken, or leaked in transit, etc. Interferences and Limitations If testing a particular sample occurs early during primary infection, no detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes A panel of Measles, Mumps, and Rubella IgG can be ordered under 1756, MMR Index Value. Reference Range Positive, Negative or Equivocal N/A	Result Indicator	Positive, Negative or Equivocal
detectable IgG may be evident. If Mumps infection is suspected, a second sample should be taken at least 14 days later. Additional Information & Notes A panel of Measles, Mumps, and Rubella IgG can be ordered under 1756, MMR Index Value. Reference Range Positive, Negative or Equivocal N/A	Unsatisfactory Specimen	specimen must be frozen at ≤-20°C and received frozen on dry ice. Specimens containing anticoagulants or preservatives. Hemolyzed, lipemic, or bacterially contaminated serum. Unlabeled or mislabeled specimens, insufficient quantity for testing, incorrect collection tube/transport media, grossly contaminated specimen, disparity between ID on sample and paperwork, improper collection, storage or transport of specimen, no test requested, test requested is not performed. If required, the absence of patient history. If required, the lack of patient history that is compatible with test requested. Test order cancelled by provider,
Additional Information & Notes A panel of Measles, Mumps, and Rubella IgG can be ordered under 1756, MMR Index Value. Reference Range Positive, Negative or Equivocal N/A	Interferences and Limitations	detectable IgG may be evident. If Mumps infection is suspected, a second
Reference Lab N/A	Additional Information & Notes	A panel of Measles, Mumps, and Rubella IgG can be ordered
· ·	Reference Range	Positive, Negative or Equivocal
Reflex testing N/A	Reference Lab	N/A
	Reflex testing	N/A