Appendix A (RFA25-001) – Cinotti Health Screening and Services Grant Program – Autism Screening Tools

Screening Population	Screening Time Interval	Recommending Organization for Screening Guidelines
All children at 18 and 24 months of age (AAP).	Routine developmental screening (all children) at 9, 18, 30 months (AAP)	American Academy of Pediatrics (AAP)
General developmental screening: All children at 9, 18, and 30 months of age (AAP).	Autism specific screening (all children) 18- 24 months (AAP)	American Academy of Child and Adolescent Psychiatry
The developmental assessment of young children and the psychiatric assessment of <u>all</u> children should routinely	Autism specific screening at any age if	( <u>AACAP</u> )
include questions about ASD symptomatology (AACAP).	triggered by concerns noted on developmental screening (AAP)	
Any adult presenting for a first-time psychiatric		
assessment (AACAP)	All young children 5 and under presenting for a developmental assessment (AACAP)	Screening Protocol Effective Dates
	All children (up to age 18) presenting for a psychiatric assessment (AACAP)	<ul><li>AAP - 2020</li><li>AACAP - 2014</li></ul>

## References:

Children: Modified Checklist for Autism in Toddlers-R/F (M-CHAT-R/F), Screening Tool for Autism in Toddlers and Young Children (STAT), and Social Communication Questionnaire (SCQ) are examples of ASD screening tools for children.

Adults: <u>Ritvo Autism Asperger's Diagnostic Scale-Revised (RAADS-R)</u> and <u>Autism Spectrum Quotient (AQ)</u> are examples of ASD screening tools for adolescents and adults.